

# Cauchy Sequences and Real Numbers

Sal Balkus

October 2020

## 1 Cauchy Sequences

A Cauchy sequence is a sequence  $x : \mathbb{N} \rightarrow \mathbb{Q}$  such that  $\forall \epsilon > 0 \exists N \in \mathbb{N}$  such that  $|x(n+p) - x(n)| < \epsilon \forall n > N$  and  $\forall p \geq 1$ . In layman's terms, the difference between a term and another term that comes after the first is arbitrarily small if the sequence goes out far enough.

One of the most useful aspects of a Cauchy sequence is that it can be used to define the set of all real numbers  $\mathbb{R}$ . A real number is simply the limit of a Cauchy sequence of rational numbers; a Cauchy sequence that converges to some limit does not necessarily have any other way to describe the number to which it converges, so we call this number "real."

**Exercise:** Prove that the sequence  $x(1) = 1, x(n) = 1 + \frac{1}{1+x(n)}$  is a Cauchy sequence.

First, we will prove the case for  $n + 1$ . Define  $\epsilon(n) = |x(n+1) - x(n)| = |1 + \frac{1}{1+x(n)} - 1 - \frac{1}{1+x(n)}| = \frac{2-x(n)^2}{(x(n)+1)^2}$  through algebraic manipulation. Then it is known that

$$\begin{aligned}\epsilon(n+1) &= \left| 1 + \frac{1}{1+x(n+1)} - x(n+1) \right| \\ &= \left| 1 + \frac{1}{1+1+\frac{1}{1+x(n)}} - 1 - \frac{1}{1+x(n)} \right| \\ &= \left| \frac{1+x(n)}{3+2x(n)} - \frac{1}{1+x(n)} \right| \\ &= \left| \frac{x(n)^2 - 2}{(3+2x(n))(1+x(n))} \right| \\ &= \left| \frac{2-x(n)^2}{x(n+1)} \right| \left| \frac{1}{3+2x(n)} \right|\end{aligned}$$

We know that  $x(1) = 1$  and the sequence is increasing therefore  $\frac{1}{3+2x(n)} < \frac{1}{5}$ . As such,  $\epsilon(n+1) < \frac{\epsilon(n)}{5} = \frac{1}{5^n}$ .

Then, we can use the telescopic sum technique to write

$$\begin{aligned} |x(n+p) - x(n)| &= |x(n+p) - x(n+p-1)| + |x(n+p-1) - x(n+p-2)| + \\ &\dots + |x(n+1) - x(n)| \\ &< \frac{1}{5^{n+p}} + \frac{1}{5^{n+p-1}} + \dots + \frac{1}{5^n} \\ &= \frac{1}{5^n} \left( \frac{1}{5^p + \frac{1}{5^{p-1}} + \dots + 1} \right) \\ &< \frac{1}{5^n} * 5 = \frac{1}{5^{n-1}} \end{aligned}$$

So, given  $\epsilon > 0$  we take  $N$  large enough such that  $\frac{1}{5^{N-1}} < \epsilon$  and this ensures that  $|x(n+p) - x(n)| < \epsilon$  for all  $n > N$  and  $p \geq 1$ , making  $x$  a Cauchy sequence.

Going further, we can also demonstrate that if a sequence is a Cauchy sequence, then that sequence is bounded, meaning that there exists  $c > 0$  such that  $|x(n)| \leq C$  for all  $n \geq 1$ .

One way to do this is a proof by contradiction. Assume that some Cauchy sequence is unbounded. Then, there does not exist  $c > 0$  such that  $|x(n)| \leq c$  for all  $n \geq 1$ . Therefore,  $\lim_{n \rightarrow \infty} x(n) = \infty$ ; the sequence diverges. If this is true then  $|x(n+p) - x(n)|$  cannot be less than some arbitrary  $\epsilon$  since  $x(n+p)$  can be infinitely large. This means that the initial assumption that  $x(n)$  is a Cauchy sequence is violated and therefore a Cauchy sequence must be bounded.